Child's Name		
Avoid Triggers: (Check all tha	at apply)	Center/staff:
□ Illness	☐ Cigarette/other smoke	☐ Food:
☐ Emotions	☐ Exercise	☐ Allergies:
☐ Weather Changes	☐ Chemical odors	☐ Other:
	_	
Green Zone: Child breathing at best Well	Yellow Zone: Child not breathing at be	est Red Zone: Danger Zone Emergency
 sleeps through the night without coughing or wheezing has no early warning signs of a asthma flare-up plays actively 	or at child care	 coughing, short of breath, wheezing neck and chest "suck in" skin between ribs, above the breastbone and collarbone when
Take Long-Term Control medications:	 has trouble doing usual activities/play, may self limit activities/ squat/hunch over decrease in appetite/difficult drinking or taking a bottle. 	
Take quick-relief medicines 15 minutes before active playtime.	Take quick–relief medicines: Adjust Long-Term Control medicines as follows unti- back in Green Zone:	col il no improvement 15 minutes after quick relief medication given and nails or lips are blue is having trouble walking or
Parent: Telephone:	Activity Restrictions:	talking cannot stop coughing
Physician:	Ozone Restrictions:	
Telephone:	Call child's parent if: child's symptoms do not improve or worsen 15 to 20	
	minutes after treatment Call the physician if:	Date:

• parent not available

's Asthma Action Plan